

This Page Is Inserted by IFW Operations
and is not a part of the Official Record

BEST AVAILABLE IMAGES

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images may include (but are not limited to):

- BLACK BORDERS
- TEXT CUT OFF AT TOP, BOTTOM OR SIDES
- FADED TEXT
- ILLEGIBLE TEXT
- SKEWED/SLANTED IMAGES
- COLORED PHOTOS
- BLACK OR VERY BLACK AND WHITE DARK PHOTOS
- GRAY SCALE DOCUMENTS

IMAGES ARE BEST AVAILABLE COPY.

As rescanning documents *will not* correct images,
please do not report the images to the
Image Problem Mailbox.




Fig. 1 Mean survival of desiccated and non-desiccated (fully-hydrated) infective juvenile *Steinernema carpocapsae* exposed to air at 25°C and 70% RH. The nematodes were desiccated through slow drying in controlled-humidity chambers following the methods of Simons and Poinar (1973). Nematode survival was determined following 24 h rehydration in water. Whiskers indicate standard error.

Fig. 2

SPECIES OF INSECT TESTED	CONDITIONING OF NEMATODES	
	DESICCATED	NON-DESICCATED
<i>Trichoplusia ni</i>	100 ± 0	100 ± 0
<i>Heliothis verescens</i>	96.6 ± 3.3	100 ± 0





Fig. 3





Fig. 4




Fig. 5




Fig. 6




Fig. 7